

Semileptonic decays of pseudoscalar mesons to scalar f_0 meson

T. M. Aliev ^{*†}, M. Savcı [‡]

Physics Department, Middle East Technical University, 06531 Ankara, Turkey

Abstract

The transition form factors of $D_s \rightarrow f_0 \ell \nu$, $D \rightarrow f_0 \ell \nu$ and $B_u \rightarrow f_0 \ell \nu$ decays are calculated in 3-point QCD sum rule method, assuming that f_0 is a quark–antiquark state with a mixture of strange and light quarks. The branching ratios of these decays are calculated in terms of the mixing angle.

^{*}e-mail: taliev@metu.edu.tr

[†]permanent address: Institute of Physics, Baku, Azerbaijan

[‡]e-mail: savci@metu.edu.tr

1 Introduction

The inner structure of the scalar mesons in terms of quarks is still an open question in particle physics and it is the subject of an intense and continuous theoretical and experimental investigations for establishing their nature (for a review, see [1]). There are numerous scenarios for the classification of the scalar mesons. The established 0^{++} mesons are divided into two groups: 1) Near and above 1 GeV , and 2) in the region $1.3\text{ GeV} \div 1.5\text{ GeV}$. The first group scalar mesons form an $SU(3)$ nonet, which contains two isosinglets, an isotriplet and two strange isodoublets. In the quark model, the flavor structure of these scalar mesons would be

$$\begin{aligned} |\sigma\rangle &= \cos\theta|\bar{n}n\rangle - \sin\theta|\bar{s}s\rangle, \\ |f_0\rangle &= \cos\theta|\bar{s}s\rangle + \sin\theta|\bar{n}n\rangle, \\ a_0^0 &= \frac{1}{\sqrt{2}}(\bar{u}u - \bar{d}d), \quad a_0^+ = u\bar{d}, \quad a_0^- = \bar{d}u, \\ \kappa^+ &= \bar{s}u, \quad \bar{\kappa}^0 = \bar{d}s, \quad \kappa^- = \bar{u}s, \end{aligned} \tag{1}$$

where $|\bar{n}n\rangle = (\bar{u}u + \bar{d}d)/\sqrt{2}$, and θ is the mixing angle. Here we take into account the fact that between isoscalars $\bar{s}s$ and $\bar{u}u + \bar{d}d$ there is mixing, which follows from experiments. Indeed the observation

$$\Gamma(J/\psi \rightarrow f_0\omega) \simeq \frac{1}{2}\Gamma(J/\psi \rightarrow f_0\phi),$$

indicates that the quark content of $f_0(980)$ is not purely $\bar{s}s$ state, but should have non-strange parts too [2]. Secondly, if $f_0(980)$ is purely $\bar{s}s$ state, then $f_0 \rightarrow \pi\pi$ should be OZI suppressed. But the decay width of $f_0(980)$ is dominated by $f_0 \rightarrow \pi\pi$ which leads to the conclusion that in $f_0(980)$ there should be $\bar{n}n$ parts as well. Therefore f_0 should be a mixture of $\bar{s}s$ and $\bar{n}n$, as is presented in Eq. (1). Analysis of the experimental data shows that the mixing angle θ lies in the range $25^\circ < \theta < 40^\circ$ and $140^\circ < \theta < 165^\circ$ [3].

Although there is another scenario where mesons below or about 1 GeV is described as a four-quark state (see for example [4]), in this work we restrict ourselves to considering the $\bar{q}q$ description for $f_0(980)$ meson, but taking into account the mixing between $\bar{s}s$ and $\bar{n}n$. In the present work we study the semileptonic decays $B^+ \rightarrow f_0(980)\ell^+\nu$, $D_{d,s}^+ \rightarrow f_0(980)\ell^+\nu$ decays in order to get information about the quark content of $f_0(980)$.

From theoretical point of view, investigation of the semileptonic decays is simpler compared to that of hadronic decays, because leptons do not participate in strong interactions. The experimental study of weak semileptonic decays of heavy flavored mesons is very important for the more accurate determination of the Cabibbo-kobayashi-Maskawa (CKM) matrix elements, their leptonic decay constants, etc.

The precise determination of the CKM matrix elements depends crucially on the possibility of controlling long distance interaction effects. So, in study of the exclusive semileptonic decays the main problem is calculation of the transition form factors, which involve the long distance QCD dynamics, belonging to the non-perturbative sector of QCD. For this reason, in calculation of the transition form factors some kind of non-perturbative

approach is needed. Among all non-perturbative approaches QCD sum rules method [5] is more powerful, since it is based on the first principles of QCD. About the most recent status of QCD sum rules, the interested readers are advised to consult [6].

Semileptonic decays $D \rightarrow \bar{K}^0 e \bar{\nu}_e$ [7], $D^+ \rightarrow K(K^{0*}) e^+ \nu_e$ [8], $D \rightarrow \pi e \bar{\nu}_e$ [9], $D \rightarrow \rho e \bar{\nu}_e$ [10], $B \rightarrow D(D^*) \ell \bar{\nu}_\ell$ [11] and $D \rightarrow \phi \ell \bar{\nu}_\ell$ [12] are all studied in the framework of 3-point QCD sum rules method.

In this work we study the semileptonic $B_u \rightarrow f_0(980) \ell^+ \nu_\ell$ and $D_{s(d)} \rightarrow f_0(980) \ell^+ \nu_\ell$ decays in the 3-point QCD sum rules method. The paper is organized as follows: In section 2, we derive the sum rules for the form factors, responsible for pseudoscalar to scalar meson transition. Section 3 is devoted to the numerical analysis of the transition form factors and discussion and contains our conclusions.

2 Pseudoscalar–scalar meson transition form factors from QCD sum rules

Pseudoscalar–scalar transition form factors are defined via the matrix element of the weak current sandwiched between initial and final meson states $\langle S(p') | \bar{q}_1 \gamma_\mu (1 - \gamma_5) q_2 | P(p) \rangle$, where q_1 and q_2 are the relevant quarks, P and S are the pseudoscalar and scalar meson states, respectively. It follows from parity conservation in strong interaction that only axial part of weak current gives non-zero contribution to this matrix element, and imposing Lorentz invariance, it can be written in terms of the form factors as follows:

$$\langle S(p') | \bar{q}_1 \gamma_\mu (1 - \gamma_5) q_2 | P(p) \rangle = -i [f_+(p + p')_\mu + f_- q_\mu] , \quad (2)$$

where $q_\mu = p_1 - p_2$.

For evaluation of these form factors in the QCD sum rule, we consider the following the 3-point correlation function

$$\Pi_\mu(p^2, p'^2, q^2) = - \int d^4x d^4y e^{i(p'y - px)} \langle 0 | T \{ J_S(y) J_\mu^A(0) J_P(x) \} | 0 \rangle , \quad (3)$$

where, $J_S = \bar{q}_2 q_2$, $J_\mu^A = \bar{q}_2 \gamma_\mu \gamma_5 q_1$ and $J_P = \bar{q}_1 \gamma_5 q_2$ are the interpolating currents of scalar and pseudoscalar mesons, and weak axial currents, respectively. It should be noted here that, $q_3 = u$, $q_2 = u$ and $q_1 = b$ for the $B_u \rightarrow f_0(980)$ transition; and $q_3 = s(d)$, $q_2 = s(d)$ and $q_1 = c$ for the $D_{s(d)} \rightarrow f_0(980)$ transition, respectively.

The decomposition of the correlation function (3) into the Lorentz structures, obviously, has the form

$$\Pi_\mu = \Pi_+(p + p')_\mu + \Pi_-(p - p')_\mu . \quad (4)$$

For the amplitudes Π_+ and Π_- , we have the following dispersion relation

$$\Pi_\pm(p^2, p'^2, Q^2) = - \frac{1}{(2\pi)^2} \int \frac{\rho_\pm(s, s', Q^2) ds ds'}{(s - p^2)(s' - p'^2)} + \text{subtraction terms} , \quad (5)$$

where $\rho_\pm$ is the corresponding spectral density and $Q^2 = -q^2 > 0$. According to QCD sum rules approach, the correlation function is calculated by the operator product expansion

(OPE) at large Euclidean momenta p^2 and p'^2 on one side, and on the other side it is calculated by inserting a complete set of intermediate states having the same quantum numbers with the currents J_S and J_P .

The phenomenological part of (3) is obtained by saturating correlator it with the lowest pseudoscalar (in our case B_u , D_s or D mesons) and scalar $f_0(980)$ mesons, yielding

$$\Pi_\mu = \frac{\langle 0 | J_S | S(p') \rangle \langle S(p') | J_\mu^A(0) | P(p) \rangle \langle P(p) | J_P(x) | 0 \rangle}{(m_S^2 - p'^2)(m_P^2 - p^2)} + \text{excited states} . \quad (6)$$

The matrix elements in Eq. (6) are defined as

$$\begin{aligned} \langle 0 | J_S | S(p') \rangle &= \lambda_S , \\ \langle P | J_P | 0 \rangle &= -i \frac{m_P^2 f_P}{m_1 + m_2} , \end{aligned} \quad (7)$$

where f_S and f_P are the leptonic decay constants of scalar and pseudoscalar mesons, and m_S and m_P are being their masses, respectively. Note that, leptonic decay constant f_S in Eq. (7) is scale dependent for which we choose the scale to be $\mu = 1 \text{ GeV}^2$, and

$$\begin{aligned} m_1 &= \begin{cases} m_b & \text{for } B_u \rightarrow f_0 \ell \nu , \\ m_c & \text{for } D_s \rightarrow f_0 \ell \nu , \quad D \rightarrow f_0 \ell \nu , \end{cases} \\ m_2 &= \begin{cases} m_u & \text{for } B_u \rightarrow f_0 \ell \nu , \quad D \rightarrow f_0 \ell \nu , \\ m_s & \text{for } D_s \rightarrow f_0 \ell \nu , \end{cases} \end{aligned}$$

Using Eqs. (2), (4), (6) and (7), for the invariant structures we get

$$\Pi_\pm = - \frac{f_P m_P^2}{m_1 + m_2} \frac{\lambda_S f_\pm}{(m_S^2 - p'^2)(m_P^2 - p^2)} . \quad (8)$$

From QCD side, the correlation function can be calculated with the help of the OPE at short distance, and in this work we will consider operators up to dimension six. The theoretical part of the correlator for the $B_s \rightarrow D_{s0}(2317) \ell \nu$ is calculated in [13], and in the present work, for the theoretical part of the corresponding sum rules, we will use the results of this work.

For the spectral densities we have

$$\Pi_+ = \frac{N_c}{4\lambda^{1/2}(s, s', Q^2)} \left[(\Delta' + \Delta) (1 + A + B) + (m_1^2 + 2m_1 m_2 + Q^2)(A + B) \right] , \quad (9)$$

$$\begin{aligned} \Pi_- &= \frac{N_c}{4\lambda^{1/2}(s, s', Q^2)} \left[(\Delta' + \Delta + m_1^2 + 2m_1 m_2 + Q^2 + 2m_1 m_2) (A - B) \right. \\ &\quad \left. + \Delta' - \Delta - 2m_1 m_2 \right] , \end{aligned} \quad (10)$$

where $N_c = 3$, $\Delta = s - m_1^2$, $\Delta' = s' - m_2^2$, and

$$A = \frac{1}{\lambda(s, s', Q^2)} \left[- (s + s' + Q^2) \Delta' + 2s' \Delta \right],$$

$$B = \frac{1}{\lambda(s, s', Q^2)} \left[- (s + s' + Q^2) \Delta' + 2s \Delta' \right].$$

For the decays under consideration, m_2 is $m_u(m_d)$ or m_s , and therefore, to take into account $SU(3)$ -violating effects, here and in all following calculations we will retain terms that linear with m_2 , while neglecting the terms higher order in m_2 .

For power corrections (PC) we get

$$\begin{aligned} \Pi_+^{PC} = & \frac{1}{2} \langle \bar{q}_2 q_2 \rangle \frac{m_1 - m_2}{rr'} + \frac{1}{4} m_2 \langle \bar{q}_2 q_2 \rangle \left(\frac{m_1^2}{r^2 r'} - \frac{2}{rr'} \right) \\ & - \frac{1}{12} m_0^2 \bar{q}_2 q_2 \left[\frac{3m_1^2(m_1 - m_2)}{r^3 r'} + \frac{2(m_1 - 2m_2)}{r r'^2} + \frac{2(2m_1 - m_2)}{r^2 r'} \right. \\ & \left. + \frac{m_1(2m_1^2 + m_1 m_2 + 2Q^2) - 2m_2(m_1^2 + Q^2)}{r^2 r'^2} \right] \\ & + \frac{4}{81} \pi \alpha_s \langle \bar{q}_2 q_2 \rangle^2 \left[- \frac{12m_1^3(m_1 - m_2)}{r^4 r'} + \frac{8m_1 m_2(m_1^2 + Q^2)}{r^2 r'^3} + \frac{56m_1 m_2}{r r'^3} \right. \\ & \left. - \frac{4m_1^2(2m_1^2 + m_1 m_2 + 2Q^2) - 8m_1 m_2(m_1^2 + Q^2)}{r^3 r'^2} - \frac{8m_1(8m_1 - 7m_2)}{r^3 r'} \right. \\ & \left. + \frac{48}{r r'^2} + \frac{48}{r^2 r'} - \frac{4(5m_1^2 - 20m_1 m_2 - 2Q^2)}{r^2 r'^2} \right] \\ & + \frac{1}{9} m_0^2 m_2 \langle \bar{q}_2 q_2 \rangle^2 \left[- \frac{m_1^2(m_1^2 + Q^2)}{r^3 r'^2} + \frac{5m_1^2 + 4Q^2}{r^2 r'^2} + \frac{6m_1^4}{r^4 r'} + \frac{10m_1^2}{r^3 r'} \right], \end{aligned} \quad (11)$$

$$\begin{aligned} \Pi_-^{PC} = & -\frac{1}{2} \langle \bar{q}_2 q_2 \rangle \frac{m_1 + m_2}{rr'} + \frac{1}{4} m_1 m_2 \langle \bar{q}_2 q_2 \rangle \left(- \frac{m_1}{r^2 r'} \right) \\ & + \frac{1}{12} m_0^2 \bar{q}_2 q_2 \left[\frac{3m_1^2(m_1 + m_2)}{r^3 r'} + \frac{2(m_1 + 3m_2)}{r r'^2} + \frac{2(3m_1 + m_2)}{r^2 r'} \right. \\ & \left. + \frac{m_1(2m_1^2 + m_1 m_2 + 2Q^2) + 2m_2(m_1^2 + Q^2)}{r^2 r'^2} \right] \\ & + \frac{1}{81} \pi \alpha_s \langle \bar{q}_2 q_2 \rangle^2 \left[\frac{12m_1^3(m_1 + m_2)}{r^4 r'} - \frac{8m_1 m_2(m_1^2 + Q^2)}{r^2 r'^3} - \frac{56m_1 m_2}{r r'^3} \right. \\ & \left. + \frac{4m_1^2(2m_1^2 + m_1 m_2 + 2Q^2) + 8m_1 m_2(m_1^2 + Q^2)}{r^3 r'^2} + \frac{8m_1(9m_1 + 7m_2)}{r^3 r'} \right. \\ & \left. + \frac{28m_1^2}{r^2 r'^2} + \frac{8}{r r'^2} - \frac{8}{r^2 r'} \right] \\ & + \frac{1}{9} m_0^2 m_2 \langle \bar{q}_2 q_2 \rangle^2 \left[\frac{m_1^2(m_1^2 + Q^2)}{r^3 r'^2} - \frac{m_1^2}{r^2 r'^2} - \frac{6m_1^4}{r^4 r'} + \frac{4}{r r'^2} - \frac{4}{r^2 r'} - \frac{24m_1^2}{r^3 r'} \right], \end{aligned} \quad (12)$$

where $r = p^2 - m_1^2$ and $r' = p'^2$. Note that the $D_s \rightarrow f_0(980)\ell^+\nu_\ell$ and $D \rightarrow f_0(980)\ell^+\nu_\ell$ decays which are considered in [14] differ from our results in three aspects:

- Our result on spectral density is two times smaller compared to that given in [14]. Since it is known that the main contribution to the sum rules comes from the spectral density, it is indispensable that our results on the form factors differ from those predicted in [14].
- In [14], part of those diagrams which are proportional to m_s are not taken into account (in our case they correspond to the terms proportional to $m_2 m_0^2 \langle \bar{q}_2 q_2 \rangle$).
- Sum rules for the form factor f_- are totally absent in [14], which could be essential for the $B_u \rightarrow f_0(980)\tau\nu_\tau$ decay.

Contribution of higher states in the physical part of the sum rules are taken into account with the help of the hadron-quark duality, i.e., corresponding spectral density for higher states is equal to the perturbative spectral density for s_0 and s'_0 starting from $s > s_0$ and $s' > s'_0$, where s and s' are the continuum thresholds in the corresponding channels.

Equating the two representations for the invariant structures $\Pi_\pm$, and applying double Borel transformation on the variables p^2 and p'^2 ($p^2 \rightarrow M^2$, $p'^2 \rightarrow M'^2$) in order to suppress the higher states and continuum contributions, we get the following sum rules for the form factors f_+ and f_- :

$$f_\pm(q^2) = -\frac{m_1 + m_2}{f_P m_P^2} \frac{1}{\lambda_S} e^{m_P^2/M^2} e^{m_S^2/M'^2} \left\{ \int ds ds' \rho_\pm(s, s', Q^2) e^{-s/M^2 - s'/M'^2} + \mathcal{B}_{M^2} \mathcal{B}_{M'^2} \Pi_\pm^{PC} \right\}. \quad (13)$$

The double Borel transformation for the quantity $1/r^n r'^m$ is defined as:

$$\mathcal{B}_{M^2} \mathcal{B}_{M'^2} \frac{1}{r^n r'^m} = (-1)^{n+m} \frac{(M^2)^{n-1}}{\Gamma(n)} \frac{(M'^2)^{m-1}}{\Gamma(m)} e^{-m_1^2/M^2}. \quad (14)$$

The integration region for the perturbative contribution is determined from the following inequalities:

$$-1 \leq \frac{2ss' + (m_1^2 - s)(s + s' + Q^2)}{\lambda^{1/2}(s, s', Q^2)(m_1^2 - s)} \leq 1. \quad (15)$$

In the calculation of the widths of the considered decays, it is necessary to know the q^2 dependence of the form factors in the whole physical region $m_\ell^2 \leq q^2 \leq q_{max}^2$.

3 Numerical analysis

In this section we present our results for the form factors $f_+(q^2)$ and $f_-(q^2)$ for the decays under consideration. The main input parameters for the sum rules are the Borel parameters

M^2 and M'^2 , continuum thresholds s_0 and s'_0 . The values of other parameters needed are: $m_b = (4.7 \pm 0.1) \text{ GeV}$ [6], $m_c = 1.4 \text{ GeV}$, $m_s = 0.15 \text{ GeV}$, $\langle \bar{u}u \rangle|_{\mu=1 \text{ GeV}} = -(0.243)^3 \text{ GeV}^3$, $\langle \bar{s}s \rangle = 0.8 \times \langle \bar{u}u \rangle$ [15]. The values of the leptonic decay constants of B_u , D_s and D mesons are determined from the analysis of the corresponding two-point correlators: $f_{B_u} = (0.14 \pm 0.01) \text{ GeV}$ [16], $f_{D_s} = (0.22 \pm 0.02) \text{ GeV}$ [17] and $f_D = (0.17 \pm 0.02) \text{ GeV}$ [6, 16, 17]. For the continuum thresholds we take the values $s_0^{B_u} = (33 \pm 2) \text{ GeV}^2$, $s_0^{D_s} = (7.7 \pm 1.1) \text{ GeV}^2$, $s_0^D = (6 \pm 0.2) \text{ GeV}^2$ and $s'_0 = 1.6 \text{ GeV}^2$ which is determined from 2-point sum rules analysis [6, 14, 16, 18].

The Borel parameters M^2 and M'^2 are the auxiliary parameters and therefore the physical quantities should be independent of them. For this reason we need to find the working regions of M^2 and M'^2 where form factors are practically independent of them.

In obtaining the working regions of M^2 and M'^2 the following two conditions should be satisfied:

- The continuum contribution should be small, and,
- power corrections should be convergent.

Our numerical analysis shows that, both conditions are satisfied in the region $10 \text{ GeV}^2 \leq M^2 \leq 20 \text{ GeV}^2$ for $B_u \rightarrow f_0 \ell \bar{\nu}_\ell$, $4 \text{ GeV}^2 \leq M^2 \leq 8 \text{ GeV}^2$ for $D_s(D) \rightarrow f_0 \ell \bar{\nu}_\ell$, and $1.2 \text{ GeV}^2 \leq M'^2 \leq 2 \text{ GeV}^2$ for all channels.

Varying the input parameters s_0 , s'_0 , f_0 , f_{D_s} , f_B and f_D in the respective regions as mentioned in the text, we get the following results for the form factors at $q^2 = 0$

$$\begin{aligned} f_+^{B_u}(0) &= 1.7(0.25 \pm 0.02) , \\ f_-^{B_u}(0) &= -1.7(0.24 \pm 0.03) , \\ f_+^D(0) &= 1.7(0.32 \pm 0.03) , \\ f_+^{D_s}(0) &= 1.7(0.27 \pm 0.02) , \end{aligned} \tag{16}$$

The multiplying factor 1.7 corresponds to the case for $\lambda_S = 0.19 \text{ GeV}^2$, and without this factor $\lambda_S = 0.35 \text{ GeV}^2$ [3]. For a comparison we present the results of the form factor f_+ for the $B_u \rightarrow f_0(980)$ transition coming from the covariant light front dynamics [19] and dispersion relation approach [20], as well as the result for the $B_u \rightarrow \pi$ transition [21].

$$\begin{aligned} f_+^{B_u \rightarrow f_0}(0) &= 0.27 \quad [19] , \\ f_+^{B_u \rightarrow f_0}(0) &= 0.09 \quad [20] , \\ f_+^{B_u \rightarrow \pi}(0) &= 0.25 \quad [21] . \end{aligned} \tag{17}$$

From a comparison of Eqs. (16) and (17) we see that, our prediction on f_+ for the $B_u \rightarrow f_0(980)$ transitions quite close to the prediction of the light front dynamics and that of $B_u \rightarrow \pi$ transition when $\lambda_S = 0.35$, and approximately three times larger compared to that of the dispersion relation approach. These close results of the form factor f_+ for the $B_u \rightarrow \pi$ and $B_u \rightarrow f_0$ transitions could indicate of the that λ_S should have the value $\lambda_S = 0.35$, which is obtained in [3] by taking $\mathcal{O}(\alpha_s)$ corrections into account.

Note that we present the form factor f_- only for the $B_u \rightarrow f_0 \tau \bar{\nu}_\tau$ decay, because this form factor can give considerable contribution to this decay.

In estimating the width of $P \rightarrow f_0(980)\ell\bar{\nu}_\ell$ decay, we need to know the q^2 dependence of the form factors $f_+(q^2)$ and $f_-(q^2)$ in the whole kinematical region $m_\ell^2 \leq q^2 \leq (m_P - m_{f_0})^2$. The q^2 dependence of the form factors can be calculated from QCD sum rules (see [8, 9]). Unfortunately QCD sum rule cannot reliably predict q^2 dependence of the form factors in the full kinematical region. The QCD sum rules can reliably predict q^2 dependence of the form factors in the region approximately 1 GeV^2 below the perturbative cut. In order to extend the dependence of the form factors on q^2 to the full kinematical region, we look such a parametrization of the form factors where it coincides with the sum rules prediction of in the above-mentioned region. Our numerical calculations shows that the best parametrization of the form factors with respect to q^2 are as follows:

$$f_P(q^2) = \frac{f_P(0)}{1 - a_P\hat{q} + b_P\hat{q}^2 - c_P\hat{q}^3 + d_P\hat{q}^4}, \quad (18)$$

where $P = B_u, D_s, D$ and $\hat{q} = q^2/m_P^2$. The values of the parameters $f_P(0)$, a_P , b_P , c_P and d_P at $\lambda_S = 0.19 \text{ GeV}^2$, are given in table 1.

	$f_+(0)$	$f_-(0)$	a	b	c	d
D_s	1.7×0.27		0.87	-0.17	0.37	1.46
D	1.7×0.32		0.89	-0.40	0.18	-1.00
B_u	1.7×0.25		0.48	-0.30	-0.47	-0.99
B_u		-1.7×0.24	0.41	-0.42	-0.95	-1.55

Table 1: Form factors for the $D_s \rightarrow f_0\ell\bar{\nu}_\ell$, $D \rightarrow f_0\ell\bar{\nu}_\ell$ and $B_u \rightarrow f_0\ell\bar{\nu}_\ell$ decays in a four-parameter fit.

The dependence of the form factors f_+ and f_- (for $B_u \rightarrow f_0\tau\bar{\nu}_\tau$ decay) are given in Figs. (1)–(4).

Using the parametrization of Eq. (2), for the $P \rightarrow f_0\ell\bar{\nu}_\ell$ differential decay width, we get

$$\begin{aligned} \frac{d\Gamma}{dq^2} = & \frac{A}{192\pi^3 m_P^3} G^2 |V_{ij}|^2 \lambda^{1/2}(m_P^2, m_{f_0}^2, q^2) \left(\frac{q^2 - m_\ell^2}{q^2} \right)^2 \\ & \times \left\{ -\frac{(2q^2 + m_\ell^2)}{2} \left[|f_+(q^2)|^2 (2m_P^2 + 2m_{f_0}^2 - q^2) + 2(m_P^2 - m_{f_0}^2) \text{Re}[f_+(q^2)f_-^*(q^2)] \right. \right. \\ & + |f_-(q^2)|^2 q^2 \left. \right] + \frac{(q^2 + 2m_\ell^2)}{q^2} \left[|f_+(q^2)|^2 (m_P^2 - m_{f_0}^2)^2 \right. \\ & \left. \left. + 2(m_P^2 - m_{f_0}^2)q^2 \text{Re}[f_+(q^2)f_-^*(q^2)] + |f_-(q^2)|^2 q^4 \right] \right\}, \quad (19) \end{aligned}$$

where

$$A = \begin{cases} \cos^2 \theta & \text{for } D_s \rightarrow f_0\ell\bar{\nu}_\ell \text{ and } , \\ \frac{\sin^2 \theta}{2} & \text{for } D \rightarrow f_0\ell\bar{\nu}_\ell \text{ and } B_u \rightarrow f_0\ell\bar{\nu}_\ell , \end{cases}$$

$$V_{ij} = \begin{cases} |V_{ub}| = 4.31 \times 10^{-3} & \text{for } B_u \rightarrow f_0 \ell \bar{\nu}_\ell , \\ |V_{cs}| = 0.96 & \text{for } D_s \rightarrow f_0 \ell \bar{\nu}_\ell , \\ |V_{cd}| = 0.23 & \text{for } D \rightarrow f_0 \ell \bar{\nu}_\ell , \end{cases}$$

Taking into account the q^2 dependence of the form factors f_+ and f_- and performing integration over q^2 and using the lifetimes of B_u , D_s and D mesons, we get the following values for the branching ratios when $\lambda_S = 0.19 \text{ GeV}^2$:

$$\begin{aligned} \mathcal{B}(B_u \rightarrow f_0 \tau \bar{\nu}_\tau) &= \frac{\sin^2 \theta}{2} \times (7.17 \times 10^{-5}) , \\ \mathcal{B}(B_u \rightarrow f_0 \mu \bar{\nu}_\mu) &= \frac{\sin^2 \theta}{2} \times (2.1 \times 10^{-4}) , \\ \mathcal{B}(B_u \rightarrow f_0 e \bar{\nu}_e) &= \frac{\sin^2 \theta}{2} \times (2.1 \times 10^{-4}) , \\ \mathcal{B}(D_s \rightarrow f_0 \mu \bar{\nu}_\mu) &= \cos^2 \theta \times (3.85 \times 10^{-3}) , \\ \mathcal{B}(D_s \rightarrow f_0 e \bar{\nu}_e) &= \cos^2 \theta \times (4.07 \times 10^{-3}) , \\ \mathcal{B}(D \rightarrow f_0 \mu \bar{\nu}_\mu) &= \frac{\sin^2 \theta}{2} \times (4.98 \times 10^{-4}) , \\ \mathcal{B}(D \rightarrow f_0 e \bar{\nu}_e) &= \frac{\sin^2 \theta}{2} \times (5.29 \times 10^{-4}) . \end{aligned} \tag{20}$$

We see from (20) that the ratios of the widths

$$\begin{aligned} R_1 &= \frac{\mathcal{B}(D \rightarrow f_0 \ell \bar{\nu}_\ell)}{\mathcal{B}(D_s \rightarrow f_0 \ell \bar{\nu}_\ell)} , \\ R_2 &= \frac{\mathcal{B}(B_u \rightarrow f_0 \ell \bar{\nu}_\ell)}{\mathcal{B}(D_s \rightarrow f_0 \ell \bar{\nu}_\ell)} , \end{aligned}$$

are directly related with the mixing angle θ . On the other hand, as far as the flavor structure of $f_0(980)$, as is given in Eq. (1), is considered, the ratio

$$R_3 = \frac{\mathcal{B}(B_u \rightarrow f_0 \ell \bar{\nu}_\ell)}{\mathcal{B}(D \rightarrow f_0 \ell \bar{\nu}_\ell)} ,$$

is independent of the mixing angle θ . Therefore, experimental measurement of the branching ratios of $B_u \rightarrow f_0 \ell \bar{\nu}_\ell$, $D_s \rightarrow f_0 \ell \bar{\nu}_\ell$ and $D \rightarrow f_0 \ell \bar{\nu}_\ell$ decays can give direct information about the mixing angle θ , as well as, about the flavor structure of $f_0(980)$ meson.

In conclusion, we study the semileptonic decay of pseudoscalar mesons to the scalar $f_0(980)$ meson. The transition form factors are calculated using 3-point QCD sum rule analysis and then we estimate the corresponding branching ratios.

References

- [1] S. Spanier and N. A. Törnqvist, "Note on scalar mesons", Phys. Lett. B **592**, 1 (2004); S. Godfrey and J. Napolitano, Rev. Mod. Phys. **71**, 1411 (1999); F. E. Close and N. A. Törnqvist, J. Phys. G **28**, 249 (2002); L. Montanet, Rep. Prog. Phys. **46**, 337 (1983); F. E. Close, Rep. Prog. Phys. **51**, 833 (1988); N. N. Achasov, Nucl. Phys. Proc. Suppl. **21**, 189 (1991); T. Barnes, prep: hep-ph/0001326.
- [2] W. -M. Yao *et al.*, Particle Data Group, J. Phys. G **33**, 1 (2006).
- [3] H. Y. Cheng, Phys. Rev. D **67**, 034024 (2004); W. Wang, Y. L. Shen, Y. Li and C. D. Lü, prep: hep-ph/0609082 (2006).
- [4] M. G. Alford and R. L. Jaffe, Nucl. Phys. B **578**, 367 (2000).
- [5] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. B **147**, 385 (1979).
- [6] P. Colangelo, A. Khodjamirian, in "At the Frontier of Particle Physics/Handbook of QCD" ed. by M. Shifman (World Scientific, Singapore, 2001), Volume **3**, 1495.
- [7] T. M. Aliev, V. L. Eletsky, Ya. I. Kogan, Sov. J. Nucl. Phys. **40**, 527 (1984).
- [8] P. Ball, V. M. Braun, H. G. Dosch, Phys. Rev. D **44**, 3567 (1991).
- [9] P. Ball, Phys. Rev. D **48**, 3190 (1993).
- [10] A. A. Ovchinnikov and V. A. Slobodenyuk, Z. Phys. C **44**, 433 (1989); V. N. Baier, and A. Grozin, Z. Phys. C **47**, 669 (1990).
- [11] A. A. Ovchinnikov, Sov. J. Nucl. Phys. **50**, 519 (1989).
- [12] D. S. Du, J. W. Li, and M. Z. Yang, Eur. Phys. J. C **37**, 173 (2004).
- [13] T. Aliev, M. Savcı, Phys. Rev. D **73**, 114010 (2006).
- [14] I. Bediaga, F. S. Navarra and M. Nielsen, Phys. Lett. B **579**, 59 (2006).
- [15] B. L. Ioffe, Prog. Part. Nucl. Phys. **56**, 232 (2006).
- [16] T. M. Aliev, V. L. Eletsky, Sov. J. Nucl. Phys. **38**, 936 (1983).
- [17] I. Bediaga and M. Nielsen, Phys. Rev. D **68**, 036001 (2003).
- [18] F. De Fazio and M. R. Pennington, Phys. Lett. B **521**, 15 (2001).
- [19] O. M. A. Leitner, B. El-Bennich, B. Loiseau and J. P. Dedonder, prep: hep-ph/0609062 (2006).
- [20] B. El-Bennich, O. M. A. Leitner, B. Loiseau and J. P. Dedonder, prep: hep-ph/0609154 (2006).
- [21] P. Ball, prep: hep-ph/0612190 (2006).

Figure captions

Fig. (1) The dependence of the form factor f_+ on q^2 at $M^2 = 15 \text{ GeV}^2$, $M'^2 = 2 \text{ GeV}^2$, $m_0^2 = 0.8 \text{ GeV}^2$, $s_0 = 33 \text{ GeV}^2$ and $s'_0 = 1.6 \text{ GeV}^2$, for the $B_u \rightarrow f_0(980)\ell\bar{\nu}_\ell$ decay.

Fig. (2) The dependence of the form factor f_- on q^2 at $M^2 = 15 \text{ GeV}^2$, $M'^2 = 2 \text{ GeV}^2$, $m_0^2 = 0.8 \text{ GeV}^2$, $s_0 = 33 \text{ GeV}^2$ and $s'_0 = 1.6 \text{ GeV}^2$, for the $B_u \rightarrow f_0(980)\ell\bar{\nu}_\ell$ decay.

Fig. (3) The dependence of the form factor f_+ on q^2 at $M^2 = 6 \text{ GeV}^2$, $M'^2 = 2 \text{ GeV}^2$, $m_0^2 = 0.6 \text{ GeV}^2$, $s_0 = 7 \text{ GeV}^2$ and $s'_0 = 1.6 \text{ GeV}^2$, for the $D_s \rightarrow f_0(980)\ell\bar{\nu}_\ell$ decay.

Fig. (4) The dependence of the form factor f_+ on q^2 at $M^2 = 6 \text{ GeV}^2$, $M'^2 = 2 \text{ GeV}^2$, $m_0^2 = 0.8 \text{ GeV}^2$, $s_0 = 6 \text{ GeV}^2$ and $s'_0 = 1.6 \text{ GeV}^2$, for the $D \rightarrow f_0(980)\ell\bar{\nu}_\ell$ decay.




Figure 1:




Figure 2:




Figure 3:




Figure 4: